

TRANSITION FROM ORDER TO CHAOS IN ROTATIONAL NUCLEI*

F.S. Stephens, M.A. Deleplanque, I.Y. Lee, A.O. Macchiavelli, D. Ward, P. Fallon, M. Cromaz, R.M. Clark, M. Descovich, R.M. Diamond, and E. Rodriguez-Vieitez
Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

Chaos in nuclei has been studied by measuring the fluctuations in energy-level spacings (the ‘nearest-neighbor distributions’ and the ‘ Δ_3 statistics’ [1]) and the results show that near the neutron binding energy the behavior is essentially chaotic [2]; whereas, near the ground state it is mainly ordered [3]. We have developed a new method to trace the behavior between these points. We study the Yb nuclei produced in heavy-ion fusion reactions, which bring high angular momentum and thermal excitation energy, E^* , into the fused system. Neutron evaporation quickly brings E^* down to about the neutron binding energy, and the angular momentum and remaining E^* are removed in a γ -ray cascade down to the ground state. This cascade covers the range between ordered and chaotic behavior in these nuclei and is the focus of our study. The Yb nuclei were chosen because nuclei in this region are deformed and exhibit rotational behavior which is essential for our method.

We have studied the narrow (valley-ridge) structure that appears in gated γ -ray spectra following heavy-ion fusion reactions and have shown that the intensity of this structure can be quantitatively related to the ratio, v/d_2 , where v is the average interaction between 2p2h states and d_2 is the average separation between such states. It has been shown that v/d_2 is, in turn, related to the chaotic behavior of a system [4]. We were able to trace the behavior from nearly fully ordered to nearly fully chaotic. It is important to note that measuring the intensity of this narrow structure does not require the separate identification of compound-damped and discrete γ rays. Both of these γ -ray types arise from events that enter and decay via the same component of the wave function and they produce similar structures in the spectrum. We use a simulation code [5] to determine the intensity of this structure, but this measurement does not rely on the use of such a code provided the statistical γ rays are removed, and the spectrum is unfolded to remove Compton-scattered γ rays.

This is a new way to explore the order-to-chaos transition in nuclei and it can be used where measuring the energy-level spacings is not possible. It looks directly at a property of the wave function rather than at level spacings. Studies of this type will give considerably more information as the simulation parameters become better established and other properties can be reliably extracted.

* This work has been supported by the U.S. DOE under Contract No. DE-AC03-76SF00098.

- [1] F.J. Dyson and M. Mehta, *J. Math. Phys.* **4**, 701 (1963).
- [2] O. Bohigas et al., *Nucl. Data for Sci. and Technology*, K.H. Böchhoff (ed.), Reidel, Dordrecht, p. 809 (1983).
- [3] J.D. Garrett et al., *Phys. Lett.* **B392**, 24 (1997).
- [4] S. Åberg, *Phys. Rev. Lett.* **64**, 3119 (1990).
- [5] F.S. Stephens et al., *Phys. Rev. Lett.* **88**, 142501, (2002); *AIP Conf. Proc.* **656**, 40 (2002).