INVESTIGATION OF ANTI-MAGNETIC ROTATION IN $^{101}\text{Pd}^*$

S. Zhu^{a,c}, U. Garg^a, A. V. Afanasjev^a, S. Frauendorf^a, B. Kharraja^a,
S. S. Ghugre^b, S. N. Chintalapudi^b, N. S. Pattabiraman^{a,b},
R. V. F. Janssens^c, M. P. Carpenter^c, F. G. Kondev^c, and T. Lauritsen^c

^a) *Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA*

^b) *IUCDAEF-Calcutta Centre, Sector III/LB-8, Bidhan Nagar, Calcutta 700 064, India*

^c) *Physics Division, Argonne National Laboratory, Argonne, IL 60439, USA*

“Magnetic rotation” has been a topic of great interest and activity in nuclear structure in the past few years. In analogy with “ferromagnetism” and “antiferromagnetism”, “antimagnetic rotation” was proposed by Frauendorf some time ago [1,2]. Experimentally, two cases in the $A\sim 100$ region have been observed and proposed as examples of “antimagnetic rotation” [3,4]. Considered as being composed of two systems of the magnetic type, the antimagnetic rotational bands appear as a regular cascade of E2 transitions in nuclei with very small deformation ($\varepsilon \leq 0.15$).

High-spin states in the nucleus ^{101}Pd have been studied with GAMMASPHERE at the ATLAS facility at ANL. The level scheme of ^{101}Pd has been considerably extended. A cascade of four “rotational-band-like” transitions was observed in this nucleus. This cascade exhibits the anticipated regular increase of transition energies with spin, and a smooth variation of the associated $\mathfrak{S}^{(1)}$ and $\mathfrak{S}^{(2)}$ moments of inertia with frequency. The configuration of the band is proposed to be $\pi(g_{9/2})_{12}^{-4} \otimes \nu(h_{11/2})_{11/2}^2 (g_{7/2}d_{5/2})_{15/2}^3$. Its antimagnetic rotational nature has been inferred from a limited lifetime measurement which was carried out in a manner similar to that used in the case of ^{100}Pd in Ref. [3]. The limits on the lifetimes are such that they indicate small deformation. This observation, combined with the rotational character and the very large estimated $\mathfrak{S}^{(2)}/B(E2)$ ratios (~ 100 times larger than those of a well-deformed collective rotor) are consistent with expectations for an antimagnetic rotational band.

* This work was supported in part by the National Science Foundation (Grant No. PHY99-01133), the U.S. Department of Energy, Office of Nuclear Physics, under Contract No. W31-109-ENG-38, and the University Grants Commission, Government of India.

- [1] S. Frauendorf, *Z. Phys.* **A385**, 163 (1997).
- [2] S. Frauendorf, *Rev. Mod. Phys.* **73**, 463 (2001).
- [3] S. Zhu et al., *Phys. Rev. C* **64**, 041302(R) (2001).
- [4] A. J. Simons et al., *Phys. Rev. Lett.* **91**, 162501 (2003).